| | | Asthma Ac | tion Plan fo | or Scho | ool C | hildr | en | Year | 20.20 | - 2021 | |
|----------------------|---|----------------------|--|--|-------------|--|--|------------|-----------------|------------|--|
| | | Name | | DOB | | Height | | jht | | | |
| • | U.V. | Personal Best Peak F | | | | | | | | | |
| | | Parent/Guardian | | | (H) | | | (C) | | | |
| | | MD/DO/NP/PAPhPh | | | | | | | | | |
| N ZONE | GO | | TAKE CONTROLLER MEDICINE EVERY DAY | | | | | | | | |
| | Green Zone (>80% of best peak flow) | | DAILY MEDICINE | | HOW MUCH | | | H | HOW OFTEN/WHEN | | |
| | No re | scue medication | | | | <u> </u> | | _ | | | |
| GREEN | | | | | | | | | | | |
| O Can | EXERCI | SE-INDUCED ASTHMA: 6 | Rescue Inhaler: | | use | | puffs | prior to h | ard exercise/ | /sports. | |
| | | (O)) a comment | ms occur with exercise, follow yellow zone. RESCUE MEDICINE | | | | | | | | |
| | T. | | | | | | | | | | |
| | Yellow ZoneTo (50-80% of best peak flow) • COUGH | | MEDICINE | HOW MUCH | | HOW OFTEN/WHEN | | | | | |
| Ę | | | Rescue medicine: | Take puffs of metered dose inhaler | | | Recheck peak flow in 15 min. If peak flow still in yellow zone, may repeat | | | | |
| | WHEEZE TIGHT CHEST | | | O Spacer | | dose once. If still in yellow zone after | | | zone after | | |
| | If still having trouble, follow red zone. | | | nebulizer treatment (s) | | | 2nd dose, call parent or doctor. STAY WITH CHILD. GIVE DRINK OF CLEAR FLUIDS. | | | | |
| | | | NOTE: Parent should contact the doctor if child needs rescue med >2 times in a week to see f a medication change is necessary. | | | | | | | | |
| RED ZONE | DANG | ER | IF IN SEVERE DI | STRESS, | CALL 9 | 11 | | 1000 | 15.50 | | |
| | RED ZONE lower than(<50% of best peak flow) • MEDICINE IS NOT HELPING • CONSTANT COUGH • WORKING HARD TO BREATHE CHILD IS VERY SICK DANGER - GET HELP! | | RESCUE MEDICINE | HOW MUCH | | HOW OFTEN/WHEN | | | | | |
| | | | Rescue medicine: | Take puffs of metered dose inhaler | | Recheck peak flow in 15 min. If peak flow still in red zone, may repeat dose once. If still in red zone after 2nd dose, call parent or doctor. | | | n. If peak | | |
| | | | | Spacer nebulizer treatment (s) | | | | | | | |
| | | | | | | | STAY WITH CHILD. GIVE DRINK OF CLEAR FLUIDS. | | | | |
| Initial | I have instructed the student in the proper use of his/her rescue medicine and the student is able to perform procedure alone and may carry inhaler. | | | I hereby release the local School Board and their agents and employees from any liability that may result from my child taking the prescribed medication. I give permission for this student to receive medications and for health care providers to exchange information regarding the care of my child. I agree to provide rescue medication to be kept at school in case of | | | | | | | |
| | /NP/PA:_ | | | emergency. | ; w provido | e rescue m | edicati | on to be k | ept at school i | in case of | |
| Date (update yearly) | | | | Parent/Guardian: | | | Date | | | | |